

TOWARDS THE COHOMOLOGY OF MODULI SPACES OF HIGHER GENUS STABLE MAPS

CLAUDIO FONTANARI

ABSTRACT. We prove that the orbifold desingularization of the moduli space of stable maps of genus $g = 1$ recently constructed by Vakil and Zinger has vanishing rational cohomology groups in odd degree $k < 10$.

1.

Let $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d)$ be the moduli space of stable maps of degree d from n -pointed curves of genus g to the projective space $\mathbb{P}^r$. For a comprehensive introduction to this fascinating geometric object we refer to the classical text [7], which not only contains a careful construction of Kontsevich moduli spaces but also outlines their crucial application to Gromov-Witten theory.

For $g = 0$ it turns out that $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^r, d)$ carries a natural structure of smooth orbifold, in particular its rational cohomology is well-defined and well-behaved. Indeed, in the last few years there has been a flurry of research about the cohomological properties of $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^r, d)$ after the first steps moved in [2]: it is worth mentioning at least the contributions [1], [3], [20], [21], [22], [16], [17], [18], [19], [9], [4], [5], [6].

For higher genus $g > 0$, instead, $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^r, d)$ can be arbitrarily singular (see [24]) and it is in general nonreduced with several components of exceptional dimension. As a consequence, a cohomological investigation of the underlying topological spaces has never been addressed and it seems to be completely out of reach. However, at least in the case $g = 1$, the beautiful construction performed in [25] has recently opened a new frontier to the research in the field.

Namely, in [25] it is shown that the closure $\overline{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d)$ of the stratum $\mathcal{M}_{1,n}^0(\mathbb{P}^r, d)$ corresponding to stable maps with smooth domain allows an orbifold desingularization

$$\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d) \rightarrow \overline{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d)$$

which can be explicitly described as a sequence of blow-ups along smooth centers. Moreover, the natural $(\mathbb{C}^*)^{r+1}$ -action on $\overline{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d)$

2000 Mathematics Subject Classification: 14H10, 14F25.

lifts to $\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d)$ and the corresponding fixed loci are explicitly determined. As a by-product, the rational cohomology groups

$$H^k(\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d), \mathbb{Q})$$

are well-defined and can be approached via localization techniques.

In the present paper we take this direction and prove the following result:

Theorem 1. *We have $H^k(\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d), \mathbb{Q}) = 0$ for every $n \geq 0$, $r \geq 1$ and odd $k < 10$.*

We point out that the natural surjective map

$$\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d) \rightarrow \overline{\mathcal{M}}_{1,n}$$

together with the well-known fact that

$$H^{11}(\overline{\mathcal{M}}_{1,11}, \mathbb{Q}) \neq 0$$

(see, for instance, [23], proof of Corollary 4.7, or [10], Proposition 2) show that in the above statement the bound on the degree k is sharp.

Our proof is based on a suitable version of the Bialynicki-Birula decomposition for singular varieties ([15] and [20]), which reduces the claim of Theorem 1 to the analogous claim on the fixed locus. Next, a careful analysis of its connected components allows us to conclude thanks to previous results on the cohomology of the moduli spaces of stable curves of genus $g \leq 1$ ([13] and [8]).

We work over the field $\mathbb{C}$ of complex numbers.

Many thanks are due to Gilberto Bini and Barbara Fantechi for stimulating conversations about this research project.

2.

We begin by recalling some more or less well-known facts about the Bialynicki-Birula decomposition for orbifolds.

Proposition 1. *Let X be a smooth projective orbifold with a $\mathbb{C}^*$ -action, let F be the fixed locus and let F_i denote its connected components. Then*

(i) *X is the disjoint union of locally closed subvarieties S_i such that every S_i retracts onto the corresponding F_i and*

$$\overline{S_i} \subseteq \bigcup_{j \geq i} S_j$$

(ii) *the Betti numbers of X are*

$$h^m(X) = \sum_i h^{m-2n_i}(F_i)$$

where n_i is the codimension of S_i .

Proof. (i) follows from [15], (1.14) and (1.15), pp. 388-389; (ii) follows from (i) and [20], Lemma 6 (i). $\square$

We also need the following technical result: please refer to [25], Section 2.1, for all non standard definitions and notation.

Lemma 1. *Fix $N \in \mathbb{N}$. Let Y be a smooth (orbi)variety and let $\{X_1, \dots, X_n\}$ be a properly intersecting collection of subvarieties of Y such that $\bigcap_{i \in I} X_i$ is smooth for every $I \subseteq \{1, \dots, n\}$ and $H^k(\bigcap_{i \in I} X_i, \mathbb{Q}) = 0$ for every odd $k < N$. Let $Y_1 := Y$ and for $i = 1, \dots, n$ let $f_i : Y_{i+1} \rightarrow Y_i$ be the blow-up of the proper transform of X_i in Y_i .*

(i) If X_i^j denotes the proper transform of X_i in Y_j , then $\bigcap_{i \in I} X_i^j$ is smooth for every $j \geq 1$ and every $I \subseteq \{j \leq i \leq n\}$ and $H^k(\bigcap_{i \in I} X_i^j, \mathbb{Q}) = 0$ for every odd $k < N$.

(ii) If moreover $H^k(Y, \mathbb{Q}) = 0$ for every odd $k < N$, then $H^k(Y_i, \mathbb{Q}) = 0$ for every odd $k < N$ and for every $i \leq n + 1$.

Proof. (i) By induction on j , the case $j = 1$ holding by assumption. We have

$$\bigcap_{i \in I} X_i^j = \bigcap_{i \in I} \text{Pr}_{X_{j-1}^{j-1}} X_i^{j-1} = \text{Pr}_{X_{j-1}^{j-1}} \bigcap_{i \in I} X_i^{j-1} = \text{Bl}_Z X$$

with $Z := \bigcap_{i \in I} X_i^{j-1} \cap X_{j-1}^{j-1}$ and $X := \bigcap_{i \in I} X_i^{j-1}$. Here the first equality holds by definition, the second one by [25], Lemma 2.3 (2), and the third one by [25], Lemma 2.3 (1).

Now, it is well-know how to compare cohomology groups after a blow-up along a smooth subvariety of an orbifold (see [11], p. 605 and Proposition on p. 606, and [14], footnote on p. 514, for the orbifold case).

In particular, it follows that if both $H^k(X, \mathbb{Q}) = 0$ and $H^k(Z, \mathbb{Q}) = 0$ for odd $k < N$, then $H^k(\text{Bl}_Z X, \mathbb{Q}) = 0$ for odd $k < N$. Hence in our case we obtain by induction both the smoothness and the vanishing result.

(ii) By induction on i , the case $i = 1$ holding by assumption. We have

$$Y_{i+1} = \text{Bl}_{X_i^i} Y_i$$

with $H^k(Y_i, \mathbb{Q}) = 0$ for odd $k < N$ by induction and $H^k(X_i^i, \mathbb{Q}) = 0$ for odd $k < N$ by (i). Hence the claim follows from the cohomological properties of blow-ups recalled above. $\square$

Proof of Theorem 1. By [25], Theorem 1.1, $\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d)$ is a smooth projective orbifold and the natural $(\mathbb{C}^*)^{r+1}$ -action on the projective space $\mathbb{P}^r$ lifts to an action on $\widetilde{\mathcal{M}}_{1,n}^0(\mathbb{P}^r, d)$. By Propostion 1 (ii), in order to prove Theorem 1 it is enough to show that the odd cohomology of the fixed loci vanishes in degree $k < 10$.

By [25], Section 1.4, such fixed loci are either

$$(1) \quad \prod \overline{\mathcal{M}}_{g,m}$$

with $g \leq 1$ and $m \in \mathbb{N}$ (see [25], p. 15, and [12], p. 541), or

$$(2) \quad \prod \overline{\mathcal{M}}_{0,i} \times \mathbb{P}^j \times \widetilde{\mathcal{M}}_{1,(I,J)}$$

with $i, j \in \mathbb{N}$ (see [25], p. 19), where $\widetilde{\mathcal{M}}_{1,(I,J)}$ is constructed in [25], pp. 26–27. Roughly speaking, $\widetilde{\mathcal{M}}_{1,(I,J)}$ is obtained from the moduli space $\overline{\mathcal{M}}_{1,I \sqcup J}$ of stable curves of genus 1 by successively blowing up certain subvarieties $\overline{\mathcal{M}}_{1,\rho}$ (where ρ is a suitable index) and their proper transforms (see [25], p. 25). We just point out a crucial fact: from the definition of $\overline{\mathcal{M}}_{1,\rho}$ (see [25], p. 23) it follows that

$$\overline{\mathcal{M}}_{1,\rho} = \overline{\mathcal{M}}_{1,h} \times \prod \overline{\mathcal{M}}_{0,k}$$

with $h, k \in \mathbb{N}$. More generally, if $\bigcap_{\rho_\alpha} \overline{\mathcal{M}}_{1,\rho_\alpha} \neq \emptyset$ for some subset of indices $\{\rho_\alpha\}$ then

$$(3) \quad \bigcap_{\rho_\alpha} \overline{\mathcal{M}}_{1,\rho_\alpha} = \overline{\mathcal{M}}_{1,s} \times \prod \overline{\mathcal{M}}_{0,t}$$

with $s, t \in \mathbb{N}$.

Now, recall that all odd degree cohomology of $\overline{\mathcal{M}}_{0,n}$ vanishes by Keel's results (see [13]) and that $H^k(\overline{\mathcal{M}}_{1,n}, \mathbb{Q}) = 0$ for odd $k < 10$ according to Getzler (see [8] and [10], eq. (4)). Hence the required vanishing trivially follows for fixed loci of type (1) and is reduced to the case of $\widetilde{\mathcal{M}}_{1,(I,J)}$ for fixed loci of type (2). On the other hand, by (3) we can apply Lemma 1 (ii) to our situation with $N = 10$, $Y = \overline{\mathcal{M}}_{1,I \sqcup J}$, $X_i = \overline{\mathcal{M}}_{1,\rho_i}$, and $Y_{n+1} = \widetilde{\mathcal{M}}_{1,(I,J)}$, so once again Getzler's claims allow us to conclude and the proof is over. $\square$

REFERENCES

- [1] K. Behrend and A. O'Halloran: On the cohomology of stable map spaces. *Invent. Math.* 154 (2003), no. 2, 385–450.
- [2] G. Bini and C. Fontanari: On the cohomology of $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^1, d)$. *Commun. Contemp. Math.* 4 (2002), no. 4, 751–761.
- [3] G. Bini and C. Fontanari: Calculating cohomology groups of $\overline{\mathcal{M}}_{0,n}(\mathbb{P}^r, d)$. *Ann. Mat. Pura Appl.* (4) 182 (2003), no. 3, 337–344.
- [4] J. A. Cox: An additive basis for the Chow ring of $\overline{\mathcal{M}}_{0,2}(P^r, 2)$. Pre-print math.AG/0501322.
- [5] J. A. Cox: A presentation for the Chow ring of $\overline{\mathcal{M}}_{0,2}(P^r, 2)$. Pre-print math.AG/0504575.
- [6] J. A. Cox: A presentation for the Chow ring $A^*(\overline{\mathcal{M}}_{0,2}(P^1, 2))$ Pre-print math.AG/0505112.

- [7] W. Fulton and R. Pandharipande: Notes on stable maps and quantum cohomology. Algebraic geometry—Santa Cruz 1995, 45–96, Proc. Sympos. Pure Math., 62, Part 2, Amer. Math. Soc., Providence, RI, 1997.
- [8] E. Getzler: The semi-classical approximation for modular operads. Comm. Math. Phys. 194 (1998), no. 2, 481–492.
- [9] E. Getzler and R. Pandharipande: The Betti numbers of $\overline{M}_{0,n}(r, d)$. J. Algebraic Geom. 15 (2006), no. 4, 709–732.
- [10] T. Graber and R. Pandharipande: Constructions of nontautological classes on moduli spaces of curves. Michigan Math. J. 51 (2003), no. 1, 93–109.
- [11] P. Griffiths and J. Harris: Principles of algebraic geometry. Pure and Applied Mathematics. Wiley-Interscience (John Wiley and Sons), New York, 1978.
- [12] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil, and E. Zaslow: Mirror symmetry. Clay Mathematics Monographs, 1. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2003.
- [13] S. Keel: Intersection theory of moduli space of stable n pointed curves of genus zero. Trans. Amer. Math. Soc. 330 (1992), no. 2, 545–574.
- [14] Y.-H. Kiem and J. Li: Desingularizations of the moduli space of rank 2 bundles over a curve. Math. Ann. 330 (2004), no. 3, 491–518.
- [15] F. Kirwan: Intersection homology and torus actions. J. Amer. Math. Soc. 1 (1988), no. 2, 385–400.
- [16] A. Mustata and M. A. Mustata: Intermediate Moduli Spaces of Stable Maps. Pre-print math.AG/0409569.
- [17] A. Mustata and M. A. Mustata: The Chow ring of $\overline{M}_{0,m}(n, d)$. Preprint math.AG/0507464.
- [18] A. Mustata and M. A. Mustata: Universal relations on stable map spaces in genus zero. Preprint math.AG/0607431.
- [19] A. Mustata and M. A. Mustata: Tautological rings of stable map spaces. Preprint math.AG/0607432.
- [20] D. Oprea: Tautological classes on the moduli spaces of stable maps to projective spaces. Pre-print math.AG/0404284.
- [21] D. Oprea: Divisors on the moduli spaces of stable maps to flag varieties and reconstruction. J. Reine Angew. Math. 586 (2005), 169–205.
- [22] D. Oprea: The tautological rings of the moduli spaces of stable maps to flag varieties. J. Algebraic Geom. 15 (2006), no. 4, 623–655.
- [23] M. Pikaart: An orbifold partition of $\overline{M}_g^n$. The moduli space of curves (Texel Island, 1994), 467–482, Progr. Math., 129, Birkhäuser Boston, Boston, MA, 1995.
- [24] R. Vakil: Murphy’s law in algebraic geometry: badly-behaved deformation spaces. Invent. Math. 164 (2006), no. 3, 569–590.
- [25] R. Vakil and A. Zinger: A Desingularization of the Main Component of the Moduli Space of Genus-One Stable Maps into $\mathbb{P}^n$. Pre-print math.AG/0603353.

E-mail address: claudio.fontanari@polito.it

Current address: Dipartimento di Matematica, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy.